


KAINUI

COCKSFOOT

A cocksfoot selected for its tiller density and its soft ryegrass-like leaves which provide it with significantly higher palatability. It is high yielding, summer-dry tolerant and has very good disease tolerance.

It is well suited for inclusion in pasture mixes or as a specialist pasture in summer-dry areas.

Drilled/broadcast with clover, and/or plantain and chicory as a specialist pasture for summer-dry prone areas at 6-8kg/ha. Alternatively, include 1-2kg/ha of Kainui in mixes with ryegrass, clovers and herbs such as plantain and chicory.



Agronomic traits

- Persistence (years): 5+ *
- Disease resistance: Very good
- Winter activity: Medium - high
- Minimum rainfall (mms): 600+ or irrigation
- Tiller density: High
- Sowing rate (kg/ha): Mixes: 2-4
Alone: 8-10

* Subject to management and climatic conditions

Animal safety

Recommended for:



Dairy



Beef



Sheep



KAINUI

COCKSFOOT

BENEFITS

- › Good tolerance of summer-dry conditions
- › Improved winter activity
- › Improved insect tolerance due to its strong root structure
- › Wont cause grass staggers or heat stress
- › Superior persistence
- › Can be sown as a pure sward
- › High quality cocksfoot for improved animal performance

