

THE MORE PALATABLE COCKSFOOT

Cropmark bred cocksfoot which was selected for leaf density and its soft ryegrass-like leaves which provide it with significantly higher palatability. It is high yielding, summer-dry tolerant and has high tiller density, and very good disease tolerance. It is well suited for inclusion in pasture mixes or as a specialist pasture in summer-dry areas.

Drilled/broadcast: with clover, and/or plantain and chicory as a specialist pasture for summer-dry prone areas at 6-8kg/ha. Alternatively, include 1-2kg/ha of Kainui in mixes with ryegrass and clovers. Suitable for all livestock types. Aerially applied: Kainui cocksfoot is available coated with Freshcote seed coating making it ideal to be mixed in with fertilizer and/or lime for aerial broadcasting over hill country. Can also be mixed with coated clover and plantain.



Agronomic traits

- Persistence (years): 5+ *
- Disease resistance: Very good
- Winter activity: Med High
- Growth peak: spring to summer
- Tiller density: High
- Sowing rate (kg/ha): Mixes: 1-2 AloneL 6-8

Benefits

- Sood tolerance of summer-dry conditions
- Improved winter activity
- Grass grub and Porina tolerant
- Won't cause grass staggers or heat stress

A New Zealand bred cocksfoot cultivar selected for strong yields combined with high palatability.



^{*}Subject to climate, insect pest pressure and management